

Quantum Phase Transition in Coupled Superconducting Quantum Dots Array with Charge Frustration

Sujit Sarkar

PoornaPrajna Institute of Scientific Research,
4 Sadashivanagar, Bangalore 5600 80, India.

(Dated: March 11, 2010)

We present the quantum phase transition in two capacitively coupled arrays of superconducting quantum dots (SQD). We consider the presence of gate voltage in each superconducting island. We show explicitly that the co-tunneling process involves with two coupled SQD arrays, near the maximum charge frustration line is not sufficient to explain the correct quantum phases with physically consistent phase boundaries. We consider another extra co-tunneling process along each chain to explain the correct quantum phases with physically consistent phase boundaries. There is no evidence of supersolid phase in our study. We use Bethe-ansatz and Abelian bosonization method to solve the problem.

Josephson junction arrays have attracted considerable interest in the recent years owing to their interesting physical properties [1–14]. Currently such arrays can be fabricated in restricted geometries both in one and two dimensions with well controlled parameters [15–18]. At the same time this is one of the paradigms to study the physics of quantum phase transition [1, 7, 8]. The Coulomb charging energy of the system is due to the small capacitance of the grains or junction that dominate strong quantum phase fluctuations of the superconducting order parameter and may drive the system into the insulating state at zero or very low temperature. The quantum fluctuations are controlled by the parameters of the system such as charging energy of the superconducting quantum dots (SQD) and Josephson coupling (E_J) between them.

In two dimensions, the universality class of these transitions has already been investigated in detail [10–14]. Nevertheless there are also quantum phase transition scenario that can take place in a Josephson junction ladder system as shown in the Fig. 1, where we consider two capacitively coupled one dimensional SQD arrays. In this simplest two-leg ladder system, we will be able to predict different quantum phases due to the interplay between Coulomb charging energy and E_J . Here we also show explicitly the role of co-tunneling effect. For large Coulomb charging energy sequential tunneling is not energetically favored process in the system. The major tunneling process in the system occurs via the co-tunneling process [19]. Its origin is quantum mechanical. In this process, tunneling occurs through the virtual state with energy equal to the on-site Coulomb charging energy. An appropriate co-tunneling process extract the correct quantum phases in the system. Otherwise, it leads to the wrong analysis of the system as we have found in Ref.[20]. In our previous studies we only emphasis the one dimensional SQD array, we only explain the importance of next-nearest-neighbor (NNN) E_J [7, 8] .

In this paper we study the quantum phase transition of capacitively coupled two arrays of SQD and they are connected through Josephson junctions. We consider the presence of gate voltage in each SQD which introduce the charge frustration in the system. At the charge frustration line the Coulomb charging energy of the system is degenerate for the difference of one Cooper pair in the island. Its also refer in the literature as a charge degeneracy point. Our main motivation is to study the different quantum phases around the maximum charge frustration line. Choi *et al.* [20] have also studied the same problem They have shown that in a coupled chain the major transport along both the chains occurs via co-tunneling of the electron-hole pairs They have not done any rigorous analytical exercise based on techniques, which are applicable in low dimensional quantum many particle systems and that seed mistakes. There are three major mistakes in that paper (1) There are two kinds of Luttinger liquid (LL) phase (describe as RL1 and RL2) but they have obtained only one. (2) Incorrect quantum phase analysis and physically inconsistent phase boundaries due to lack of correct analytical derivations and physical interpretations. (3). There is no evidence of super solid (SS) phase in this model.

We show explicitly in this study that a single co-tunneling process in the two leg SQD ladder system is not sufficient to produce the correct quantum phases with physically consistent phase boundaries. We also show that the co-tunneling process along each chain is also necessary for the correct phase boundaries. In our model system, superconducting islands are connected with E_J . The charging energies along each array are the on-site Coulomb charging energy due to the self capacitance, $E_0 = \frac{e^2}{2C_0}$ and the junction charging energy, $E_1 = \frac{e^2}{2C_1}$ due to junction capacitance C_1 . The arrays are coupled to each other via the capacitance C_I , it contributes charging energy, $E_I = \frac{e^2}{2C_I}$. But there are no Josephson coupling between the two arrays. The array capacitances are small, which




FIG. 1: This is the schematic diagram of our model systems. Two arrays of superconducting quantum dots are connected through the capacitance, C_I . C_0 is the self capacitance of the dot and C_1 is the junction capacitance. In our superconducting circuit, small circles denote the dots and crosses denote the Josephson junctions. V_g is the gate voltage applied in each dot.

leads to $E_J \ll E_0, E_1$. Therefore each arrays are in the insulating phase in the absence of co-tunneling effect. Here we consider the limit $E_I \ll E_0, E_1$. The applied gate voltage induce a charge in every superconducting island by an amount, $n_g = \frac{C_0 V_g}{2e}$, this applied gate voltage introduce the charge frustration in the system. We do the quantum phase analysis near the maximum charge frustration line ($|n_g - N - 1/2| \ll 1$), around this line we obtain several interesting quantum phases which have not been noticed in the previous studies [5, 6, 20]. The Hamiltonian of the system is given by

$$H = 2e^2 \sum_{l', xx'} (n_l(x) - n_g) C_{l,l'}^{-1}(x, x') (n_l(x') - n_g) - E_J \sum_{l,x} \cos(\theta_l(x) - \theta_l(x+1)) \quad (1)$$

$n_l(x)$ is the number of Cooper pairs and $\theta_l(x)$ is the phase of the superconducting order parameter at the site x of the l th array. $n_l(x)$ and $\theta_l(x)$ are the quantum mechanically conjugate variables. One can write the capacitance matrix in the block form.

$$C_{l,l'}^{-1}(x, x') = C(x, x')A + \delta_{x,x'} C_I B \quad (2)$$

Where

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$B = \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}$$

$$C(x, x') = C_0 \delta_{x,x'} + C_1 [2\delta_{x,x'} - \delta_{x,x'+1} - \delta_{x,x'-1}] \quad (3)$$

$$H = H_c^{(0)} + H_c^{(1)} + H_J \quad (4)$$

with the components

$$H_c^{(0)} = U_0 \sum_x [n_+(x) - 2n_g]^2 + V_0 \sum_x (n_-(x))^2 \quad (5)$$

$$H_c^{(1)} = U_1 \sum_x [n_+(x) - 2n_g][n_+(x+1) - 2n_g] + V_1 \sum_x n_-(x)n_-(x+1) \quad (6)$$

$$H_J = -E_J \sum_{l,x} \cos(\theta_l(x) - \theta_l(x+1)) \quad (7)$$

Where $n_{\pm}(x) = n_1(x) \pm n_2(x)$ and the coupling strengths are given by $U \sim 2E_0$, $U_1 = 4\frac{C_1}{C_0}$, $V_0 \sim E_I$ and $V_1 \sim \frac{C_1 C_I}{E_I}$. The first term of the Hamiltonian $H_c^{(0)}$ contains an important message regarding the difference between the charging energy at near to the maximum charge frustration line and the particle-hole symmetric line ($n_g = 0$). The charge configurations which do not satisfy the condition for $n_+(x) = 1$ at the maximum charge frustration line generates a gap in the excitation spectrum of the order of on site charging energy. The ground state of $H_c^{(0)}$ separated from the excited state by the gap of the order of E_I . This excited state has two fold degeneracy for every values of x , corresponding to $n_-(x) = \pm 1$. It is convenient to work in the charge configuration with $n_+(x) = 0$ and $n_-(x) = \pm 1$ is termed as a reduce Hilbert space of the problem. Presence of finite E_J lifts this degeneracy and the ground state of $H_c^{(0)}$ is mixed with the state $n_-(x) = \pm 2$. Now the relevant reduce Hilbert space, $n_+(x) = 0$ and $n_-(x) = 0, \pm 2$. Here we mention the effective model of our system, wherein we follow the references. They have found the effective Hamiltonian up to the second order in $\frac{E_J}{E_0}$.

$$H_{eff} = P[H + H_J \frac{1-P}{E - H_C^0} H_J]P \quad (8)$$

Here P is the projection operator onto the reduce space and finally they have obtained the effective Hamiltonian.

$$H_{eff} = \gamma J \sum_x S^z(x) S^z(x+1) - \frac{J}{2} \sum_x [S^+(x) S^-(x+1) + h.c.] \quad (9)$$

where the XY component of exchange interaction and the z component of exchange anisotropy are respectively $J = \frac{E_J^2}{4E_0}$ and $\gamma = \frac{16\lambda^2 E_J^2}{E_J^2}$. No analytical expression for λ^2 is given in Ref. . Here we estimate $\lambda^2 = \frac{C_1 E_0}{C_I E_I}$. Pseudo spin operators of the above Hamiltonian are the following:

$$\begin{aligned} S^z(x) &= P \frac{n_1(x) - n_2(x)}{2} P \\ S^+(x) &= P e^{-i\theta_1(x)} (1 - P) e^{-i\theta_2(x)} P \\ S^-(x) &= P e^{-i\theta_2(x)} (1 - P) e^{-i\theta_1(x)} P. \end{aligned} \quad (10)$$

Now we would like to explain the pseudo-spin terms of the Hamiltonian in terms of charge representation from where we start. The first term of the Hamiltonian originates from the first term of Eq. 9 which the Coulomb charging energy of the SQD in the form of $n(x)$. The second term of the Hamiltonian originates from the co-tunneling process, i.e, from the second term of Eq. 9. They have obtained the Hamiltonian in an effective one dimensional Hamiltonian but they have not done any correct analytical calculations based on the Abelian bosonization and Bethe ansatz methods which are suitable for one dimensional quantum many body system. Here we would like to do the quantum phase boundaries analysis based on the exact Bethe ansatz calculation based results. We will present the short comings of this effective Hamiltonian from that analysis.

$$\begin{aligned} S^z(x) &= \psi^\dagger(x)\psi(x) - 1/2, \\ S^-(x) &= \psi(x) \exp[i\pi \sum_{j=-\infty}^{x-1} n_j], \\ S^+(x) &= \psi^\dagger(x) \exp[-i\pi \sum_{j=-\infty}^{x-1} n_j], \end{aligned} \quad (11)$$

where $n(x) = \psi^\dagger(x)\psi(x)$ is the fermion number at the site x .

$$\begin{aligned} H_{eff} &= -\gamma E_J \sum_x (\psi^\dagger(x)\psi(x) - 1/2) \\ &\quad (\psi^\dagger(x+1)\psi(x+1) - 1/2) \\ &\quad - \frac{J}{2} \sum_x (\psi^\dagger(x+1)\psi(x) + \text{h.c.}) \end{aligned} \quad (12)$$

In order to study the continuum field theory of these Hamiltonians, we recast the spinless fermion operators in terms of field operators by this relation [21] $\psi(x) = [e^{ik_F x} \psi_R(x) + e^{-ik_F x} \psi_L(x)]$ where $\psi_R(x)$ and $\psi_L(x)$ describe the second-quantized fields of right- and the left-moving fermions respectively. We want to express the fermionic fields in terms of bosonic fields by the relation $\psi_r(x) = \frac{U_r}{\sqrt{2\pi\alpha}} e^{-i(r\phi(x) - \theta(x))}$, where r is denoting the chirality of the fermionic fields, right (1) or left movers (-1). The operators U_r commute with the bosonic field. U_r of different species commute and U_r of the same species anti commute. ϕ field corresponds to the quantum (bosonic) fluctuations of spin and θ is the dual field of ϕ . They are related by the relation $\phi_R = \theta - \phi$ and $\phi_L = \theta + \phi$. After doing the continuum field theory exercise, H_{eff} become

$$H_{eff} = H_0 + \frac{\gamma J}{2\pi\alpha^2} \int dx : \cos(4\sqrt{K_1}\phi(x)) : \quad (13)$$




FIG. 2: We present the one dimensional schematic diagram for quantum phases as a function of Luttinger liquid parameters (K). In the figure, we present also values of K at the phase boundaries. The value of K at the wings of the arrows denote the value of K at the phase boundaries. CDW is the charge density wave phase. RL2 is the repulsive Luttinger liquid phase of second kind. SC is the superconducting phase. The limit $K = 2$ only achieve when the system is in the first order commensurability, i.e., each dots are occupied with integer number of Cooper pairs. MI is the Mott-insulating phase of the system. The presence of gate voltage is not explicit in K , therefore the RL1 phase not appears in figure, we can only achieve this phase when the applied gate voltage exceed the CDW gap.

$H_0 = \frac{v}{2\pi} \int dx [(\partial_x \theta)^2 + (\partial_x \phi)^2]$ is the non-interacting part of the Hamiltonian. The collective velocity of the system (v) and K_1 are the two LL parameter. We use exact Bethe ansatz solution to calculate

$$K_1 = \frac{\pi}{\pi + 2\sin^{-1}\gamma} \quad (14)$$

For $J < \gamma J$ with relatively small applied gate voltage, the antiferromagnetic Ising interaction dominates the physics of anisotropic Heisenberg chain. When the applied gate voltage is large the chain is in the ferromagnetic state. In the language of interacting Cooper pairs, the Neel phase is the Charge density wave phase with period 2 i.e there is only one Cooper pair in every two sites. For the ferromagnetic phase system is in the Mott insulating phase. The emergence of two LL phases can be ascribed to the following reasons: RL1 and RL2, respectively occur due to the commensurate-incommensurate transition and the criticality of the Heisenberg XY model. The system is either in RL1 for $K < 1/2$ or in RL2 for $K > 1/2$. The physical significance of RL1 is that the coupling term is relevant but the applied gate voltage on SC island breaks the CDW gapped phase, whereas in RL2, system is gapless for $K_1 > 1/2$.

In Fig. 2, we present quantum phases of our study as a function of K . We apply the Luther-Emery (LE) trick [22] in the massive phase of sine-Gordon field theory to evaluate $K_1 (=1/4)$ at the phase boundary between the CDW and RL1 and also for CDW and RL2. RL1 phase is not explicit in the figure because we only achieve this phase under the application of gate voltage when it exceed the gap of CDW state. The value of K_1 at the phase boundary between RL2 and CDW phase is $1/4$. From the analysis of K_1 we obtain $16\lambda^2 E_I^2 = -E_J^2$, this condition is unphysical because λ , E_I and E_J are (+ve)

quantities. The value of K is 1 at the phase boundary between the RL2 and superconducting phase. The analysis yields the condition $16\lambda^2 E_I^2 = 0$, which is again unphysical. There is no chance to get superconductivity in this model Hamiltonian. We surprise that without doing any correct quantum analysis of this model Hamiltonian, the authors of Ref. [20] have claimed the existence of SC. To get the correct physical behavior between the different quantum phases, one has to be considered the Co-tunneling processes along each chain separately. We have realized during our calculations that to get an attractive interaction between the Jordan-Wigner (spinless) fermions, we have to consider the higher order expansion in $\frac{E_J}{E_{C0}}$. This higher order expansion leads to the virtual state with energies exceeding E_{C0} . In this second order process, extra contribution appears as

$$H_{extra} = \frac{-3E_J^2}{4E_0} \sum_x S^z(x) S^z(x+1) - \frac{E_J^2}{E_0} \sum_x (S^\dagger(x+2) S^-(x) + h.c) \quad (15)$$

[6–8, 23]. The total effective Hamiltonian of the system under the combined co-tunneling process is

$$H_{eff} = (\gamma J - \frac{3E_J^2}{4E_0}) \sum_x S^z(x) S^z(x+1) - \frac{J}{2} \sum_x (S^\dagger(x) S^-(x+1) + h.c) - \frac{E_J^2}{E_0} \sum_x (S^\dagger(x+2) S^-(x) + h.c) \quad (16)$$

After doing the quantum field theory, we get the effective Hamiltonian

$$H_{eff} = H_0 + \frac{\gamma J - \frac{3E_J^2}{4E_0}}{2\pi\alpha^2} \int dx : \cos(4\sqrt{K_2}\phi(x)) : \quad (17)$$

LL parameter of H_{eff} is

$$K_2 = \sqrt{\frac{J}{J + 4/\pi(\gamma J - \frac{3E_J^2}{4E_0})}} \quad (18)$$

As we notice from the Fig. 2 that value of $K_2 = 1/4$ at the phase boundary between RL2 and CDW phase. The parametric condition at the phase boundary is the following $\frac{16\lambda^2 E_I^2}{E_J^2} = \frac{15\pi}{4} + 3$. This condition is consistent physically. Therefore we prove the importance of this extra co-tunneling process. We see from the Fig. 2 that value of the K_2 is 1 at the phase boundary between the RL2 and superconducting phase. The parametric analysis for this value of K_2 yields $16\lambda^2 E_I^2 = 3E_J^2$, we obtain, $E_J^2 = 8\lambda^2 E_I^2$. This is a physically realizable condition for phase boundaries. It is clear from the Fig. 2 that there is no simultaneous presence of CDW phase

and SC phase. SC phase occurs only when $K > 1$ and CDW phase occurs only when $1/2 < K < 1$. Therefore there is no evidence of SS phase for this model system.

Conclusions: We have found the all correct quantum phases of this model Hamiltonian based on the rigorous analytical calculations. We obtain the condition for physically consistent phase boundaries which were absent in the previous studies. There is no evidence of super solid phase.

Acknowledgment: The author would like to acknowledge CCMT of the Physics Department of IISc for extended facility and also Prof. R. Srikanth for reading the manuscript.

-
- [1] S. L. Sondhi *et al.*, Rev. Mod. Phys. **69**, 315 (1997).
 - [2] R. Fazio, and H. van der Zant, Physics Report **355**, 235 (2001).
 - [3] L. J. Geerligs *et al.*, Phys. Rev. Lett. **63**, 326 (1989).
 - [4] H. S. J. van der Zant *et al.*, Euro. Phys. Lett. **119**, 541 (1992).
 - [5] R. M. Bradley, and S. Doniach, Phys. Rev. B **30**, 1138 (1984).
 - [6] L. I. Glazmann, and A. I. Larkin, Phys. Rev. Lett. **39**, 3786 (1997).
 - [7] S. Sarkar, Euro. Phys. Lett, **71** 980 (2005).
 - [8] S. Sarkar, Phys. Rev. B **75**, 014528 (2007).
 - [9] R. Baltin, and K. H. Wagenblast, Euro. Phys. Lett, **39**, 7 (1997).
 - [10] E. Granto and J. M. Kosterlitz, Phys. Rev. Lett. **65**, 1267 (1990).
 - [11] E. Granto, Phys. Rev. B **42**, 4797 (1990).
 - [12] E. Orignac and T. Giamarchi, Phys. Rev. B **64**, 144515 (2001).
 - [13] H. Lee and M. C. Cha, Phys. Rev. B **65**, 172505 (2002).
 - [14] M. P. A. Fisher, G. Grinstein and S. M. Girvin, Phys. Rev. Lett **64**, 587 (1990).
 - [15] W. Kuo and C. D. Chen, Phys. Rev. Lett. **87**, 186804 (2001).
 - [16] H. M. Jaeger *et al.* Phys. Rev. B **40**, 182 (1989).
 - [17] C. D. Chen *et al.*, Phys. Rev. B **51**, 15645 (1995).
 - [18] E. Chow, P. Delsing, and D. B. Haviland, Phys. Rev. Lett. **81**, 204 (1998).
 - [19] H. Bruus, and K. Flensberg, in *Many Body Quantum Theory in Condensed Matter Physics* (Oxford University Press, New York 2004).
 - [20] Mahn-Soo Choi *et al.*, Phys. Rev. Lett **81**, 4240 (1998).
 - [21] T. Giamarchi, in *Quantum Physics in One Dimension* (Clarendon Press, Oxford 2004).
 - [22] We follow the Luther-Emery trick during the analysis. One can write the sine-Gordon Hamiltonian for arbitrary commensurability as $H_2 = H_{01} + \lambda \int dx \cos(2n\sqrt{K}\phi(x))$, where n is the commensurability and λ is the coupling strength. H_{01} is the free part of the Hamiltonian. We know that for the spinless fermions, $\psi_R^\dagger \psi_L + \psi_L^\dagger \psi_R = \frac{1}{2\pi\alpha^2} \int dx \cos(2\sqrt{K}\phi(x))$, which is similar to the analytical expression of sine-Gordon coupling term but with the wrong coefficient inside the cosine. One can set $\phi(x) = 2\sqrt{K}\phi(x)$ then the above

equation become $H_3 = H_{01} + \lambda \int dx \cos(2\tilde{\phi}(x))$. K and $\tilde{K}$ are related by the relation, $K = \frac{\tilde{K}}{n^2}$. At the phase boundary, $\tilde{K} = 1$ that implies $K = 1/n^2$.

[23] D. Giuliano, and P. Sodano, Nucl. Phys. B **711**, 480 (2005).